



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 1/113 // C12N 15/51</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/35709</b> <b>(43) International Publication Date:</b> 14 November 1996 (14.11.96)
<b>(21) International Application Number:</b> PCT/US96/06388 <b>(22) International Filing Date:</b> 9 May 1996 (09.05.96)  <b>(30) Priority Data:</b> 08/439,680 12 May 1995 (12.05.95) US 08/571,643 13 December 1995 (13.12.95) US  <b>(71) Applicant:</b> SCHERING CORPORATION [US/US]; 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).  <b>(72) Inventors:</b> RAMANATHAN, Lata; 32 Valley Way, West Orange, NJ 07052 (US). WENDEL, Michele; 7 Robin Road, Fanwood, NJ 07023 (US).  <b>(74) Agents:</b> LUNN, Paul, G. et al.; Schering-Plough Corporation, Patent Dept. K-6-1 1990, 2000 Galloping Hill Road, Kenilworth, NJ 07033-0530 (US).	<b>(81) Designated States:</b> AL, AM, AU, AZ, BB, BG, BR, BY, CA, CN, CZ, EE, GE, HU, IS, JP, KG, KR, KZ, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> METHOD FOR REFOLDING INSOLUBLE AGGREGATES OF HEPATITIS C VIRUS PROTEASE  <b>(57) Abstract</b>  A method for solubilizing and refolding insoluble aggregates of HCV protease.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**METHOD FOR REFOLDING INSOLUBLE AGGREGATES OF**  
**HEPATITIS C VIRUS PROTEASE**

5

**BACKGROUND OF THE INVENTION**

10

Hepatitis C virus (HCV) is considered to be the major etiological agent of non-A non-B (NANB) hepatitis, chronic liver disease, and hepatocellular carcinoma (HCC) around the world. The viral infection accounts for greater than 90% of transfusion -associated hepatitis in U.S. and it is the predominant form of hepatitis in adults over 40 years of age. Almost all of the infections result in chronic hepatitis and nearly 20% develop liver cirrhosis.

The virus particle has not been identified due to the lack of an efficient *in vitro* replication system and the extremely low amount of HCV particles in infected liver tissues or blood. However, molecular cloning of the viral genome has been accomplished by isolating the messenger RNA (mRNA) from the serum of infected chimpanzees then cloned using recombinant methodologies. [Grakoui A. *et al.* J. Virol. 67: 1385 - 1395 (1993)] It is now known that HCV contains a positive strand RNA genome comprising approximately 9400 nucleotides, whose organization is similar to that of flaviviruses and pestiviruses. The genome of HCV, like that of flavi- and pestiviruses, encodes a single large polyprotein of about 3000 amino acids which undergoes proteolysis to form mature viral proteins in infected cells.

20  
25  
30

Cell-free translation of the viral polyprotein and cell culture expression studies have established that the HCV polyprotein is processed by cellular and viral proteases to produce the putative structural and nonstructural (NS) proteins. At least nine mature viral proteins are produced from the polyprotein by specific proteolysis. The order and nomenclature of the cleavage products are as follows: NH<sub>2</sub>-C-E1-E2-NS2-NS3-NS4A-NS4B-NS5A-NS5B-COOH.(Fig 1). The three

35

amino terminal putative structural proteins, C (capsid), E1, and E2 ( two envelope glycoproteins), are believed to be cleaved by host signal peptidases of the endoplasmic reticulum(ER) . The host enzyme is also responsible for generating the amino terminus of NS2 . The proteolytic processing of the nonstructural proteins are carried out by the viral proteases: NS2-3 and NS3, contained within the viral polyprotein. The NS2-3 protease catalyzes the cleavage between NS2 and NS3. It is a metalloprotease and requires both NS2 and the protease domain of NS3. The NS3 protease catalyzes the rest of the cleavages the substrates in the nonstructural part of the polyprotein. The NS3 protein contains 631 amino acid residues and is comprised of two enzymatic domains: the protease domain contained within amino acid residues 1-181 and a helicase ATPase domain contained within the rest of the protein. It is not known if the 70 kD NS3 protein is cleaved further in infected cells to separate the protease domain from the helicase domain, however, no cleavage has been observed in cell culture expression studies.

The NS3 protease is a member of the serine class of enzymes. It contains His, Asp, and Ser as the catalytic triad, Ser being the active site residue. Mutation of the Ser residue abolishes the cleavages at substrates NS3/4A, NS4A/4B, NS4B/5A, and NS5A/5B. The cleavage between NS3 and NS4A is intramolecular, whereas the cleavages at NS 4A/4B, 4B/5A, 5A/5B sites occur in *trans* .

Experiments using transient expression of various forms of HCV NS polyproteins in mammalian cells have established that the NS3 serine protease is necessary but not sufficient for efficient processing of all these cleavages. Like flaviviruses, the HCV NS3 protease also requires a cofactor to catalyze some of these cleavage reactions. In addition to the serine protease NS3, the NS4A protein is absolutely required for the cleavage of the substrate at the 4B/5A site and increases the efficiency of cleavage of the substrate between 5A/5B, and possibly 4A/4B.

Because the HCV NS3 protease cleaves the non-structural HCV proteins which are necessary for the HCV replication, the NS3 protease can be a target for the development of therapeutic agents against the HCV virus. The gene encoding the HCV NS3 protein has been cloned

- 3 -

as disclosed in U.S. Patent No. 5,371,017, however, not in a soluble active form. If the HCV protease is to be useful as a target in a screen to discover therapeutic agents, the protease must be produced in a soluble active form. Thus, there is a need for a soluble active form of the HCV protease which can be produced in large quantities to be used in high throughput screen to detect inhibitors of the protease and for structural studies. We have cloned and expressed the catalytic domain of NS3 protease as a native protein and as fusion proteins in *E. coli* and in Yeast. Fusion tags were used to facilitate purification and secretion into periplasmic space. All of these constructions resulted in expression of NS3 protein only in insoluble form. Various attempts which include growing bacteria in different media and temperatures, expressing in different strains of *E. coli* failed to produce expression of soluble NS3. Thus, there is a need for a soluble active form of the HCV protease which can be used in a screen to test for potential therapeutic agents.

#### Summary Of The Invention

The present invention fills this need by providing for a process for producing soluble, proteolytically active, refolded HCV protease from insoluble, bacterially produced HCV protease aggregates. Insoluble, aggregates of HCV NS3 protease are extracted from bacteria producing said aggregates. The aggregates of protease are then solubilized in a buffer containing a denaturing reagent. The solubilized protease from are then placed in a buffer containing a reducing agent said buffer having an acidic pH. The denaturing reagent is then removed from the buffer under conditions wherein the buffer maintains an acidic pH. The pH of the buffer containing the protease is then raised in a stepwise manner to a pH of about 7 - 8 so as to produce properly refolded soluble, active NS3 protease.

In a preferred embodiment of the present invention, the insoluble protease is first extracted from the bacteria by homogenization or sonication of the bacteria. The aggregates containing the bacteria are then solubilized in a 5 M solution of guanidine hydrochloride (GuHCl). The NS3 protease is then purified from high molecular weight aggregates by size exclusion chromatography, as for example by applying the solution to a SEPHACRYL S-300 size exclusion gel chromatography.

Fractions containing the NS3 protease are collected and the solution comprised of 5 M solution of GuHCl is diluted to about 0.1 M GuHCl in a refolding buffer containing dithiothreitol and lauryl maltoside. The diluted solution is then applied to a reverse phase chromatography column and pools containing the NS3 protease collected. The pH of the protease fractions is then raised in a stepwise manner to about 7.4 - 7.8 so as to produce properly refolded soluble, active NS3 protease.

#### Brief Description Of The Figures

10

Figure 1 schematically depicts the HCV polyprotein.

Figure 2 depicts the recombinant synthesis of plasmid pBJ1015.

15 

Figure 3 depicts the recombinant synthesis of plasmid pTS56-9.

Figure 4 depicts the recombinant synthesis of plasmid pT5His/HIV/183.

#### Detailed Description Of The Invention

20

The teachings of all references cited are incorporated herein in their entirety by reference.

25 

The amino acid sequence of the NS3 protease catalytic domain is shown in SEQ ID NO: 1. Prior to the present invention the NS3 protease could not be produced in a soluble form in sufficient quantities for extraction and purification. The present invention provides for a method to solubilize and refold bacterially produced soluble HCV protease.

30

35 

According to the present invention, soluble HCV NS3 protease can be produced having the sequences shown in SEQ ID NO: 1 and SEQ ID NO: 4. The NS3 protease can also have a histidine tag fused to its amino acid terminus for use in purifying the protein on a nickel ( $\text{Ni}^{2+}$ ) coated resin. See SEQ ID NO: 5. The protease is produced as insoluble aggregates, i.e. inclusion bodies, in bacteria such as *E. coli*.



The insoluble HCV NS3 protease is first extracted from the bacteria by homogenization or sonication of the bacteria. The aggregates containing the bacteria are then solubilized in a solubilizing agent. Suitable solubilizing agents are guanidine hydrochloride (GuHCl), urea and glutathiocyanate. Preferably the solubilizing agent is a 5 M solution of GuHCl. In a preferred embodiment, the solubilized NS3 protease is then purified from high molecular weight aggregates by size exclusion chromatography, as for example by applying the solution to a SEPHACRYL S-300 size exclusion gel column. Fractions containing the NS3 protease in the solubilizing agent are diluted in a refolding buffer containing a reducing agent. Examples of suitable reducing agents are dithiothreitol (DTT), dithioerythritol (DET) and  $\beta$ -mercaptoethanol. The preferred refolding buffer contains about 10% DTT. The refolding buffer also preferably contains a non-ionic detergent. Examples of non-ionic detergents are lauryl maltoside, a polyoxyethylene ether such as TRITON X-100<sup>®</sup>, Nonidet P-40<sup>®</sup>, a polyoxyethylene 9 -lauryl ether such as THESIT<sup>®</sup>, (3-[(3-Cholamidopropyl)-dimethylammonio]-1-propanesulfonate) (CHAPS), and octylglucoside. Preferably the insoluble aggregates of protease are solubilized in 5M GuHCl. Purified fractions from a size exclusion gel column are pooled and diluted to about 0.1 M GuHCl in a refolding buffer comprised of 10% dithiothreitol and 0.1% lauryl maltoside. The diluted solution is then applied to a reverse phase chromatography column and pools containing the NS3 protease collected. The pH of the protease fractions is then raised in a stepwise manner to about 7 - 8, preferably 7.4 - 7.8, so as to produce properly refolded soluble, active NS3 protease.

DNA encoding the NS3 protease of this invention can be prepared by chemical synthesis using the known nucleic acid sequence [Ratner *et al.*, Nucleic Acids Res. 13:5007 (1985)] and standard methods such as the phosphoramidite solid support method of Matteucci *et al.* [J Am. Chem. Soc. 103:3185 (1981)] or the method of Yoo *et al.* [J. Biol. Chem. 264:17078 (1989)]. See also Glick, Bernard R. and Pasternak, *Molecular Biotechnology* : pages 55 - 63, (ASM Press, Washington, D.C. 1994). The gene encoding the protease can also be obtained using the plasmid disclosed in Grakoui, A., Wychofski, C., Lin, C., Feinstone, S. M., and Rice, C. M., Expression and Identification of Hepatitis C Virus polyprotein Cleavage

- 6 -

Products, *J. Virol* 67:1385-1395 (1993). Also, the nucleic acid encoding HCV protease can be isolated, amplified and cloned (from patients infected with the HCV virus). Furthermore, the HCV genome has been disclosed in PCT WO 89/04669 and are available from the  
5 American Type Culture Collection (ATCC), 12301 Parklawn Drive, Rockville, MD under ATCC accession no. 40394.

Of course, because of the degeneracy of the genetic code, there are many functionally equivalent nucleic acid sequences that can encode mature human HCV protease as defined herein. Such  
10 functionally equivalent sequences, which can readily be prepared using known methods such as chemical synthesis, PCR employing modified primers and site-directed mutagenesis, are within the scope of this invention.

15 As used herein, the term "transformed bacteria" means bacteria that have been genetically engineered to produce a mammalian protein. Such genetic engineering usually entails the introduction of an expression vector into a bacterium. The expression vector is capable of autonomous replication and protein expression relative to genes in the  
20 bacterial genome. Construction of bacterial expression is well known in the art, provided the nucleotide sequence encoding a desired protein is known or otherwise available. For example, DeBoer in U.S. Pat. No. 4,551,433 discloses promoters for use in bacterial expression vectors; Goeddel *et al.* in U.S. Pat. No. 4,601,980 and Riggs, in U.S. Pat. No.  
25 4,431,739 disclose the production of mammalian proteins by *E. coli* expression systems; and Riggs *supra*, Ferretti *et al.* *Proc. Natl. Acad. Sci.* 83:599 (1986), Sproat *et al.*, *Nucleic Acid Research* 13:2959 (1985) and Mullenbach *et al.*, *J. Biol. Chem* 261:719 (1986) disclose how to construct  
30 synthetic genes for expression in bacteria. Many bacterial expression vectors are available commercially and through the American Type Culture Collection (ATCC), Rockville, Maryland.

Insertion of DNA encoding human HCV protease into a vector is easily accomplished when the termini of both the DNA and  
35 the vector comprise the same restriction site. If this is not the case, it may be necessary to modify the termini of the DNA and/or vector by



digesting back single-stranded DNA overhangs generated by restriction endonuclease cleavage to produce blunt ends, or to achieve the same result by filling in the single-stranded termini with an appropriate DNA polymerase. Alternatively, any site desired may be produced by ligating nucleotide sequences (linkers) onto the termini. Such linkers may comprise specific oligonucleotide sequences that define desired restriction sites. The cleaved vector and the DNA fragments may also be modified if required by homopolymeric tailing.

Many *E. coli*-compatible expression vectors can be used to produce soluble HCV NS3 protease of the present invention, including but not limited to vectors containing bacterial or bacteriophage promoters such as the *Tac*, *Lac*, *Trp*, *Lac UV5*,  $l P_r$  and  $l P_L$  promoters. Preferably, a vector selected will have expression control sequences that permit regulation of the rate of HCV protease expression. Then, HCV protease production can be regulated to avoid overproduction that could prove toxic to the host cells. Most preferred is a vector comprising, from 5' to 3' (upstream to downstream), a *Tac* promoter, a *lac Iq* repressor gene and DNA encoding mature human HCV protease. The vectors chosen for use in this invention may also encode secretory leaders such as the ompA or protein A leader, as long as such leaders are cleaved during post-translational processing to produce mature HCV protease or if the leaders are not cleaved, the leaders do not interfere with the enzymatic activity of the protease.

Fusion peptides will typically be made by either recombinant nucleic acid methods or by synthetic polypeptide methods. Techniques for nucleic acid manipulation and expression are described generally, e.g., in Sambrook, et al. (1989) *Molecular Cloning: A Laboratory Manual* (2d ed.), vols. 1-3, Cold Spring Harbor Laboratory; and Ausubel, et al. (eds.) (1993) *Current Protocols in Molecular Biology*, Greene and Wiley, NY. Techniques for synthesis of polypeptides are described, e.g., in Merrifield (1963) *J. Amer. Chem. Soc.* 85:2149-2156; Merrifield (1986) *Science* 232: 341-347; and Stewart et al (1984)., "Solid Phase Peptide Synthesis" (2nd Edition), Pierce Chemical Co., Rockford, IL.; and Atherton, et al. (1989) *Solid Phase Peptide Synthesis: A Practical*

*Approach*, IRL Press, Oxford; and Grant (1992) *Synthetic Peptides: A User's Guide*, W.H. Freeman, NY.

The smaller peptides such as the NS4A cofactor, SEQ ID NO: 6, 7  
5 and 8, and the substrates 5A/5B, SEQ ID NO: 5, and 4B/5A, SEQ ID NO: 9  
can be synthesized by a suitable method such as by exclusive solid phase  
synthesis, partial solid phase methods, fragment condensation or  
classical solution synthesis. The polypeptides are preferably prepared by  
solid phase peptide synthesis as described by Merrifield, J. Am. Chem.  
10 Soc. 85:2149 (1963). The synthesis is carried out with amino acids that  
are protected at the alpha-amino terminus. Trifunctional amino acids  
with labile side-chains are also protected with suitable groups to prevent  
undesired chemical reactions from occurring during the assembly of the  
polypeptides. The alpha-amino protecting group is selectively removed  
15 to allow subsequent reaction to take place at the amino-terminus. The  
conditions for the removal of the alpha-amino protecting group do not  
remove the side-chain protecting groups.

The alpha-amino protecting groups are those known to be useful  
20 in the art of stepwise polypeptide synthesis. Included are acyl type  
protecting groups (e.g., formyl, trifluoroacetyl, acetyl), aryl type  
protecting groups (e.g., biotinyl), aromatic urethane type protecting  
groups [e.g., benzyloxycarbonyl (Cbz), substituted benzyloxycarbonyl and  
9-fluorenylmethyloxy-carbonyl (Fmoc)], aliphatic urethane protecting  
25 groups [e.g., t-butyloxycarbonyl (tBoc), isopropyloxycarbonyl,  
cyclohexyloxycarbonyl] and alkyl type protecting groups (e.g., benzyl,  
triphenylmethyl). The preferred protecting groups are tBoc and Fmoc,  
thus the peptides are said to be synthesized by tBoc and Fmoc chemistry,  
respectively.

30

The side-chain protecting groups selected must remain intact  
during coupling and not be removed during the deprotection of the  
amino-terminus protecting group or during coupling conditions.  
The side-chain protecting groups must also be removable upon the  
35 completion of synthesis, using reaction conditions that will not alter  
the finished polypeptide. In tBoc chemistry, the side-chain protecting  
groups for trifunctional amino acids are mostly benzyl based. In  
Fmoc chemistry, they are mostly tert.-butyl or trityl based.

In tBoc chemistry, the preferred side-chain protecting groups are tosyl for Arg, cyclohexyl for Asp, 4-methylbenzyl (and acetamidomethyl) for Cys, benzyl for Glu, Ser and Thr, 5 benzyloxymethyl (and dinitrophenyl) for His, 2-Cl-benzyloxycarbonyl for Lys, formyl for Trp and 2-bromobenzyl for Tyr. In Fmoc chemistry, the preferred side-chain protecting groups are 2,2,5,7,8-pentamethylchroman-6-sulfonyl (Pmc) or 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl (Pbf) for Arg, trityl for 10 Asn, Cys, Gln and His, tert. butyl for Asp, Glu, Ser, Thr and Tyr, tBoc for Lys and Trp.

For the synthesis of phosphopeptides, either direct or post-assembly incorporation of the phosphate group is used. In the direct 15 incorporation strategy, the phosphate group on Ser, Thr or Tyr may be protected by methyl, benzyl or tert.butyl in Fmoc chemistry or by methyl, benzyl or phenyl in tBoc chemistry. Direct incorporation of phosphotyrosine without phosphate protection can also be used in Fmoc chemistry. In the post-assembly incorporation strategy, the 20 unprotected hydroxyl group of Ser, Thr or Tyr was derivatized on solid phase with di-tert.butyl-, dibenzyl- or dimethyl-N,N'-diisopropylphosphoramidite and then oxidized by tert.butylhydroperoxide.

25 Solid phase synthesis is usually carried out from the carboxyl-terminus by coupling the alpha-amino protected (side-chain protected) amino acid to a suitable solid support. An ester linkage is formed when the attachment is made to a chloromethyl, chlortrityl or hydroxymethyl resin, and the resulting polypeptide will have a 30 free carboxyl group at the C-terminus. Alternatively, when an amide resin such as benzhydrylamine or p-methylbenzhydrylamine resin (for tBoc chemistry) and Rink amide or PAL resin (for Fmoc chemistry) is used, an amide bond is formed and the resulting polypeptide will have a carboxamide group at the C-terminus. These 35 resins, whether polystyrene- or polyamide-based or polyethyleneglycol-grafted, with or without a handle or linker, with or without the first amino acid attached, are commercially available, and their preparations have been described by Stewart et al (1984).

"Solid Phase Peptide Synthesis" (2nd Edition), Pierce Chemical Co., Rockford, IL.; and Bayer & Rapp (1986) *Chem. Pept. Prot.* 3, 3; and Atherton, et al. (1989) *Solid Phase Peptide Synthesis: A Practical Approach*, IRL Press, Oxford.

5

The C-terminal amino acid, protected at the side-chain if necessary and at the alpha-amino group, is attached to a hydroxymethyl resin using various activating agents including dicyclohexylcarbodiimide (DCC), N,N'-diisopropylcarbodiimide  
10 DIPC DI) and carbonyldiimidazole (CDI). It can be attached to chloromethyl or chlorotriyl resin directly in its cesium tetramethylammonium salt form or in the presence of triethylamine (TEA) or diisopropylethylamine (DIEA). First amino acid attachment to an amide resin is the same as amide bond formation  
15 during coupling reactions

Following the attachment to the resin support, the alpha-amino protecting group is removed using various reagents depending on the protecting chemistry (e.g., tBoc, Fmoc). The extent  
20 of Fmoc removal can be monitored at 300-320 nm or by a conductivity cell. After removal of the alpha-amino protecting group, the remaining protected amino acids are coupled stepwise in the required order to obtain the desired sequence.

25 Various activating agents can be used for the coupling reactions including DCC, DIPC DI, 2-chloro-1,3-dimethylimidium hexafluorophosphate (CIP), benzotriazol-1-yl-oxy-tris-(dimethylamino)-phosphonium hexafluorophosphate (BOP) and its pyrrolidine analog (PyBOP), bromo-tris-pyrrolidino-phosphonium  
30 hexafluorophosphate (PyBroP), O -(benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and its tetrafluoroborate analog (TBTU) or its pyrrolidine analog (HBPyU), O -(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) and its tetrafluoroborate analog  
35 (TATU) or pyrrolidine analog (HAPyU). The most common catalytic additives used in coupling reactions include 4-dimethylaminopyridine (DMAP), 3-hydroxy-3,4-dihydro-4-oxo-1,2,3-benzotriazine (HODhbt), N-hydroxybenzotriazole (HOBt) and 1-

hydroxy-7-azabenzotriazole (HOAt). Each protected amino acid is used in excess (>2.0 equivalents), and the couplings are usually carried out in N-methylpyrrolidone (NMP) or in DMF, CH<sub>2</sub>Cl<sub>2</sub> or mixtures thereof. The extent of completion of the coupling reaction can be monitored at each stage, e.g., by the ninhydrin reaction as described by Kaiser *et al.*, *Anal. Biochem.* 34:595 (1970). In cases where incomplete coupling is found, the coupling reaction is extended and repeated and may have chaotropic salts added. The coupling reactions can be performed automatically with commercially available instruments such as ABI model 430A, 431A and 433A peptide synthesizers.

After the entire assembly of the desired polypeptide, the polypeptide-resin is cleaved with a reagent with proper scavengers. The Fmoc peptides are usually cleaved and deprotected by TFA with scavengers (e.g., H<sub>2</sub>O, ethanedithiol, phenol and thioanisole). The tBoc peptides are usually cleaved and deprotected with liquid HF for 1-2 hours at -5 to 0°C, which cleaves the polypeptide from the resin and removes most of the side-chain protecting groups. Scavengers such as anisole, dimethylsulfide and p-thiocresol are usually used with the liquid HF to prevent cations formed during the cleavage from alkylating and acylating the amino acid residues present in the polypeptide. The formyl group of Trp and dinitrophenyl group of His need to be removed, respectively, by piperidine and thiophenol in DMF prior to the HF cleavage. The acetamidomethyl group of Cys can be removed by mercury(II) acetate and alternatively by iodine, thallium (III) trifluoroacetate or silver tetrafluoroborate which simultaneously oxidize cysteine to cystine. Other strong acids used for tBoc peptide cleavage and deprotection include trifluoromethanesulfonic acid (TFMSA) and trimethylsilyltrifluoroacetate (TMSOTf).

The following examples are included to illustrate the present invention but not to limit it.

### Example 1

#### Production of HCV NS3 Protease

5

##### A. Plasmid constructions.

Several plasmids were designed and constructed using standard recombinant DNA techniques (Sambrook, Fritsch & Maniatis) to express the HCV protease in *E. coli* (Fig 2-7). All HCV specific sequences originated from the parental plasmid pBRTM/HCV 1-3011 (Grakoui *et al.* 1993). To express the N-terminal 183 amino acid versions of the protease, a stop codon was inserted into the HCV genome using synthetic oligonucleotides (Fig. 3). The plasmids designed to express the N-terminal 246 amino acid residues were generated by the natural NcoI restriction site at the C-terminus.

- i) Construction of the plasmid pBJ1015 (Figure 2)
- 20 The plasmid pBRTM/HCV 1-3011 containing the entire HCV genome (Grakoui A., *et al.*, *J. Virol.* 67: 1385-1395) was digested with the restriction enzymes Sca I and Hpa I and the 7138 bp (base pair) DNA fragment was isolated and cloned to the Sma I site of pSP72 (Promega) to produce the plasmid, pRJ201. The plasmid pRJ 201 was digested with
- 25 Msc I and the 2106 bp Msc I fragment was isolated and cloned into the Sma I site of the plasmid pBD7. The resulting plasmid pMBM48 was digested with Kas I and Nco I, and the 734 bp DNA fragment after blunt ending with Klenow polymerase was isolated and cloned into Nco I digested, klenow polymerase treated pTrc HIS B seq expression plasmid
- 30 (Invitrogen). The ligation regenerated a Nco I site at the 5' end and Nsi I site at the 3' end of HCV sequence. The plasmid pTHB HCV NS3 was then digested with Nco I and Nsi I, and treated with klenow polymerase and T4 DNA polymerase, to produce a blunt ended 738 bp DNA fragment which was isolated and cloned into Asp I cut, klenow
- 35 polymerase treated expression plasmid pQE30 (HIV). The resulting plasmid pBJ 1015 expresses HCV NS3 (246 amino acids) protease.



(ii) Construction of the plasmid pTS 56-9 with a stop codon after amino acid 183 (Figure 3)

5 The plasmid pTHB HCV NS3 was digested with Nco I, treated with klenow polymerase, then digested with Bst Y I; and the DNA fragment containing HCV sequence was isolated and cloned into Sma I and Bgl II digested pSP72. The resulting plasmid pTS 49-27 was then digested with Bgl II and Hpa I and ligated with a double stranded oligonucleotide:

10 GA TCA CCG GTC TAG ATCT

T GGC CAG ATC TAGA (SEQ ID NO 3) to produce pTS 56-9.

Thus, a stop codon was placed directly at the end of DNA encoding the protease catalytic domain of the NS3 protein. This enabled the HCV protease to be expressed independently from the helicase domain of the  
15 NS3 protein.

(iii) Construction of the plasmid pT5 His HIV-NS3 (Figure 4)

20 The plasmid pTS56-9 was digested with Bgl II, and treated with Klenow polymerase to fill in 5' ends. The plasmid was then digested with NgoM I and the blunt ended Bgl II/NgoMI fragment containing the NS3 sequence was isolated and ligated to the SglI, Klenow treated NgmMI cut and Sal I klenowed pBJ 1015. The resulting plasmid is  
25 designated pT5His HIV 183.

## Example 2

### Refolding of Insoluble HCV NS3 Protease

30

The present example describes a novel process for the refolding of HCV NS3 protease which does not have a solubilizing motif from an *E. coli* inclusion body pellet. This procedure can be used to generate purified enzyme for activity assays and structural studies.

35

Extraction and Purification of His-HIV 183 from the *E. coli* inclusion body pellet

*E. coli* cells harboring the plasmid for HisHIV183 was used to transform a culture of *E. coli* strain M15 [pREP] (Qiagen), which over-expresses the *lac* repressor, according to methods recommended by commercial source. M15 [pREP] bacteria harboring recombinant plasmids were grown overnight in 20-10-5 broth supplemented with 100µg/ml ampicillin and 25µg/ml kanamycin. Cultures were diluted to O.D.600 of 0.1, then grown at 37°C to O.D.600 of 0.6 to 0.8, after which IPTG was added to a final concentration of 1mM. At post-induction 2 to 3 hours, the cells were harvested by pelleting, and the cell pellets were washed with 100mM Tris, pH 7.5. were pelleted by centrifugation. The cell pellet was resuspended in 10 ml of 0.1M Tris-HCl, 5mM EDTA, pH 8.0 (Buffer A) for each gm wet weight of pellet. The pellet was homogenized and resuspended using a Dounce homogenizer. The suspension was clarified by centrifugation at 20,000 x g for 30 minutes at 4°C. The pellet was sequentially washed with the following five buffers:

1. Buffer A
2. 1.0M sodium chloride (NaCl) in buffer A
3. 1.0% Triton X-100 in buffer A
4. Buffer A
5. 1.0 M Guanidine HCl ( GuHCl) in buffer A.

The washed pellet was solubilized with 5M GuHCl, 1% beta mercaptoethanol in buffer A (3 ml per gm wet wt. of pellet) using a Dounce homogenizer and centrifuged at 100,000 x g for 30 minutes at 4°C. Purification of denatured HisHIV183 from high molecular weight aggregates was accomplished by size exclusion on a SEPHACRYL S-300 gel filtration column.

In particular, an 8 ml sample of the 5.0M GuHCl *E. coli* extract was applied to a 160 ml Pharmacia S-300 column (1.6 x 100 cm) at a flow rate of 1.0 ml/min. The column buffer was comprised of 5.0 M GuHCl, 0.1 M Tris-HCl, pH 8.0, and 5.0 mM EDTA. The fraction size was 5.0 ml. Appropriate fractions were pooled based on the results of SDS-PAGE, as well as N-terminal sequence analysis of the protein transferred to a Pro-Blot.

Detergent-assisted refolding of HCV-protease

The protein was concentrated by ultrafiltration using a 43 mm  
5 Amicon YM10 membrane to 1.0 mg per ml in 5M GuHCl, 0.1M Tris-HCl  
pH 8.0, 1.0 mM EDTA, 1.0% beta-mercaptoethanol. It was then diluted  
50-fold to 0.1M GuHCl in refolding buffer (100 mM sodium phosphate  
pH 8.0, 10mM DTT, 0.1% lauryl maltoside) and the mixture was  
10 incubated on ice for at least one hour. A 25 ml sample containing 500 µg  
of the protein in the refolding buffer was applied to a Pro-RPC HR 3/5  
reversed phase chromatography column. The applied sample contained  
500 µg protein in 25 ml of refolding buffer. To the column was then  
applied a solution B comprised of 99.9% H<sub>2</sub>O + 0.1% trifluoroacetic acid  
(TFA). A 10 ml volume of solution C [10% H<sub>2</sub>O, 90% acetonitrile (AcN)  
15 + 0.1% TFA] was applied to the column at a 0 - 60% gradient into  
solution B at a flow rate of 0.5ml/min. and a fraction size of 0.5ml. The  
fractions were monitored at A<sub>214</sub>; 2.0 absorbance units full scale (AUFS).

Fractions containing the protein (corresponding to peak 1) were  
20 pooled for renaturation by stepwise dialysis. The fractions were first  
dialysed in 0.1% TFA in 25% glycerol overnight at 4°C. These pooled  
fractions had a concentration of 0.1% TFA, 40% acetonitrile and a pH of  
less than 1. The fractions were then dialyzed in 0.01% TFA in 25%  
glycerol overnight at 4°C raising the pH to about 2; then dialyzed in  
25 0.001% TFA in 25% glycerol for 3.0 hours raising the pH to about 3; then  
dialyzed for 3 hours at 4°C in 50 mM NaPO<sub>4</sub>, pH 6.0, 10 mM DTT in 25%  
glycerol raising the pH to about 6. The protein was then dialyzed for 3.0  
hours at 4°C in 50 mM NaPO<sub>4</sub>, pH 7.0, 0.15 M NaCl, 10 mM DTT in 25%  
glycerol; and then finally dialyzed in 50 mM NaPO<sub>4</sub>, pH 7.8, 0.3 M NaCl,  
30 10 mM DTT, 0.2% Tween 20 in 25% glycerol. This resulted in purified,  
refolded, soluble, active HCV NS3 protease resulting in a solution  
having a pH of about 7.4 - 7.8.

Far UV circular dichroism (CD) analysis of the protein was used  
35 to monitor the refolding from an acid denatured state to a folded state at  
neutral pH. The protein recovery was monitored by a UV scan and SDS-  
PAGE analysis.

Results:Detergent-assisted Refolding of His-HIV183

5 HisHIV183 was quantitatively extracted from an *E. coli* inclusion body pellet. SDS-PAGE analysis at the various stages of extraction shows that sequential washes are essential to remove significant amounts of the contaminating proteins. HisHIV183 was extracted from the washed inclusion body pellet in the presence of 5M GuHCl. The 5M GuHCl  
10 extract was applied to a SEPHACRYL S-300 column and the appropriate fractions were pooled based on SDS-PAGE analysis. The amino acid sequence of the first ten residues was verified.

Refolding was performed at very low concentrations of protein,  
15 in the presence of DTT, lauryl maltoside and glycerol at 4°C. The diluted protein was concentrated on a Pro-RPC reversed phase column. Two peaks were obtained based on the UV and protein profile. Only Peak 1 has yielded soluble protein after stepwise dialysis. Far UV CD spectral analysis was used to monitor refolding from a denatured state at acid pH  
20 to a folded state at neutral pH. At pH 7.4, the protein was found to exhibit significant amounts of secondary structure that is consistent with that of beta sheet protein. At low pH, the CD spectrum showed that it is fully random coil, having a minimal molar ellipticity at 200nm. The ratio of this minimum at 200nm to that of the shoulder at 220 nm is  
25 approximately 4:1. This ratio decreased when the secondary structure formation occurred at neutral pH.

A UV scan at each step of dialysis showed that the protein recovery was >90% up to pH 7.0 and that there was no light scattering  
30 effect due to protein aggregates. SDS-PAGE analysis also indicated that there was no loss of protein up to pH 7.0 during refolding. Precipitation of protein occurred at the last step of dialysis, and the soluble protein was clarified by centrifugation. The overall protein recovery was about 10%. The refolded protein was found to be active in a trans-cleavage  
35 assay using the *in vitro*-translated 5A/5B substrate in the presence of 4A peptide.

### Example 3

#### Analysis of NS3 Protease Activity By *In Vitro* Translation Assay

5        To detect HCV NS3 protease activity in *trans*, we have expressed a 40 kD protein containing the NS5A/5B cleavage site in cell-free translation system and used that as the substrate for the enzyme. The substrate protein produces two protein products of apparent molecular weight 12.5 kD (NS 5A') and 27 kD (NS5B') upon cleavage by the HCV  
10    NS3 protease.

      The plasmid pTS102 encoding the substrate 5A/5B was linearized by digestion with EcoR I and was transcribed using T7 RNA polymerase *in vitro*. The RNA was translated in presence of <sup>35</sup>S methionine in  
15    rabbit reticulocyte lysates according to the manufacturer's (Promega ) protocol to produce HCV specific protein. In a 20 µl total reaction mixture containing 10mM Tris, pH 7.5, 1mM DTT, 0.5mM EDTA, and 10% glycerol was placed 2 to 8 µl of <sup>35</sup>S methionine-labeled translated 5A/5B substrate. The reaction was started with the addition of 10µl of  
20    HCV NS3 protease (SEQ ID NO: 2) with an approximately equimolar amount (2 µM) of the carboxyterminal 33 mer cofactor NS4A (SEQ ID NO: 7) in solubilization buffer (50mM Na Phosphate, pH 7.8, 0.3M NaCl, 0.2% Tween 20, 10 mM DTT or BME, 10% glycerol), and incubated at 30°C for about one hour. Reactions were stopped by adding an equal  
25    volume of 2X Laemmli sample buffer (Enprotech Inc.) and heating at 100°C for 3 minutes. Reaction products were separated by SDS PAGE electrophoresis; gels were fixed, dried and subjected to autoradiography.

      The assay was able to cleave 5A/5B substrate in a dose responsive  
30    manner, producing the expected cleaved products: 5A (12.5 kD) and 5B (27 kD) as shown by SDS PAGE analysis. The production of cleaved 5A and 5B polypeptides from the 5A/5B substrate is proof that soluble, active, refolded HCV protease was indeed produced by the process of the present invention.

Example 4Detergent-Assisted Refolding of the Catalytic Domain His-HIV 183

HCV protease catalytic domain has been expressed in high  
5 concentrations in *E.coli* as an inclusion body pellet. and is amenable to  
refolding studies. His-HIVNS3183 contains a six-residue polyhistidine  
tag, a 27 residue HIV protease cleavage sequence and a serine protease  
domain of 183 amino acids. His-HIVNS3 183 was extracted with 5M  
10 GuHCl according to the procedure of Example 2. A sample of the 5.0M  
GuHCl *E. coli* extract was applied to a 500 ml Pharmacia S-300 column  
(5.0 x 100 cm) at a flow rate of 4ml/min. The column buffer was the  
same buffer used in Example 2. About 100mg of highly purified protein  
was obtained for refolding studies.

15 The fractions containing the protein (1.0 mg / ml) were collected  
and diluted 50-fold in buffer A (100 mM sodium phosphate pH 7.8, 25%  
glycerol, 0.1% lauryl maltoside and 10 mM DTT) and immediately  
applied to a POROS 20R1 reversed phase column. A main peak and a  
shoulder were eluted with a 0-60% acetonitrile gradient in 0.1 %TFA.  
20 Only the main peak, not the shoulder, yielded active protease using a  
stepwise dialysis procedure.

Fractions containing the protein (corresponding to peak 1) were  
pooled for renaturation by stepwise dialysis. The fractions were first  
25 dialysed in 0.1% TFA in 25% glycerol overnight at 4°C. These pooled  
fractions had a concentration of 0.1% TFA, 40% acetonitrile and a pH of  
less than 1. The fractions were then dialyzed in 0.01% TFA in 25%  
glycerol overnight at 4°C raising the pH to about 2; then dialyzed in  
0.001% TFA in 25% glycerol for 3.0 hours raising the pH to about 3; then  
30 dialyzed for 3 hours at 4°C in 50 mM NaPO<sub>4</sub>, pH 6.0, 10 mM DTT in 25%  
glycerol raising the pH to about 6. The protein was then dialyzed for 3.0  
hours at 4°C in 50 mM NaPO<sub>4</sub>, pH 7.0, 0.15 M NaCl, 10 mM DTT in 25%  
glycerol; and then finally dialyzed in 50 mM NaPO<sub>4</sub>, pH 7.8, 0.3 M NaCl,  
10 mM DTT, in 25% glycerol. This resulted in purified, refolded,  
35 soluble, active HCV NS3 protease resulting in a solution having a pH of  
about 7.4 - 7.8. This resulted in an approximate 27% yield of active  
protease (>95% purity).



- 19 -

The refolded protein was found to be active in the presence of a NS4A peptide (33-mer) in the *in vitro*-translation assay using a truncated 5A-5B substrate. Three small scale refolding experiments (1.0 & 10.0 and 20 mg ) gave reproducible yields (30%) of active soluble protease. We performed a loading study on the reverse phase column to improve the recovery of refolded protein. A 2.5 mg scale refolding gave 27% recovery of active protease. Refolding of HCV protease from a one liter fermentation is estimated to give 4-5 mg of active protein.

10 We have studied the enhancement activity of NS4A peptides on the activity of refolded HCV protease in the SPA assay. Kinetics of this enzyme has been determined with the unlabeled peptide in the HPLC assay. (Table)

15

Table Kinetics of Refolded HCV Protease Catalytic Domain  
Determined in the presence of NS4A (22-54)

20 Non-Linear Regression

$K_m = 63.626 \pm 19.834 \mu M$

$V_{max} = 22.9 \pm 3.397 \text{ pmoles / min / } 0.5 \mu g \text{ enzyme}$

$k_{cat} = 1.05 \text{ min}^{-1}$

25  $k_{cat} / K_m = 264.7 \text{ M}^{-1} \text{s}^{-1}$

### Preliminary Detergent-Assisted Refolding of NS3 631

30 The full-length HCV protease NS3 631 was extracted from an *E. coli* inclusion body pellet and purified using Sephacryl S-300 chromatography. Forty milligrams of highly purified NS3 631 has been obtained from a six liter fermentation. This protein migrated as a doublet on SDS-PAGE under reducing conditions. N-terminal  
35 sequencing of the two immunoreactive bands indicated that the majority of the protein has a blocked N-terminus. The biochemical basis for the heterogeneity is unknown. Using modified detergent-assisted refolding scheme that was described for HisHIV183, low amounts of

soluble protein was obtained. The procedure was modified by including 0.5M arginine hydrochloride in the refolding buffer. The refolded protein showed activity in the presence of NS4A peptide in the *in vitro*-translation assay using truncated 5A-5B as a substrate.

5

10

15

20

25

30

35

## SEQUENCE LISTING

5

## (1) GENERAL INFORMATION:

(i) APPLICANT: Schering Corporation

10

(ii) TITLE OF INVENTION: Method for Refolding Insoluble  
Aggregates of Hepatitis C Virus Protease

(iii) NUMBER OF SEQUENCES: 9

15

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Schering-Plough Corporation

(B) STREET: 2000 Galloping Hill Road

(C) CITY: Kenilworth

20

(D) STATE: New Jersey

(E) COUNTRY: USA

(F) ZIP: 07033-0530

## (v) COMPUTER READABLE FORM:

25

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: Apple Macintosh

(C) OPERATING SYSTEM: Macintosh 7.1

(D) SOFTWARE: Microsoft Word 5.1a

30

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:

(B) FILING DATE:

(C) CLASSIFICATION:

35

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/439,680

(B) FILING DATE: May 12, 1995

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/571,643

(B) FILING DATE: December 13, 1995

(viii) ATTORNEY/AGENT INFORMATION:

- 5 (A) NAME: Lunn, Paul G.  
(B) REGISTRATION NUMBER: 32,743  
(C) REFERENCE/DOCKET NUMBER: JB0508K

(ix) TELECOMMUNICATION INFORMATION:

- 10 (A) TELEPHONE: 908-298-5061  
(B) TELEFAX: 908-298-5388

(2) INFORMATION FOR SEQ ID NO:1:

15 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 549 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

- (A) NAME/KEY: HCV NS3 Protease

25

30 GCG CCC ATC ACG GCG TAC GCC CAG CAG ACG AGA GGC CTC CTA GGG 45  
Ala Pro Ile Thr Ala Tyr Ala Gln Gln Thr Arg Gly Leu Leu Gly  
1 5 10 15  
TGT ATA ATC ACC AGC CTG ACT GGC CGG GAC AAA AAC CAA GTG GAG 90  
Cys Ile Ile Thr Ser Leu Thr Gly Arg Asp Lys Asn Gln Val Glu  
20 25 30  
35 GGT GAG GTC CAG ATC GTG TCA ACT GCT ACC CAA ACC TTC CTG GCA 135  
Gly Glu Val Gln Ile Val Ser Thr Ala Thr Gln Thr Phe Leu Ala  
35 40 45

ACG TGC ATC AAT GGG GTA TGC TGG ACT GTC TAC CAC GGG GCC GGA 180  
 Thr Cys Ile Asn Gly Val Cys Trp Thr Val Tyr His Gly Ala Gly  
 50 55 60

5  
 ACG AGG ACC ATC GCA TCA CCC AAG GGT CCT GTC ATC CAG ATG TAT 225  
 Thr Arg Thr Ile Ala Ser Pro Lys Gly Pro Val Ile Gln Met Tyr  
 65 70 75

10  
 ACC AAT GTG GAC CAA GAC CTT GTG GGC TGG CCC GCT CCT CAA GGT 270  
 Thr Asn Val Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly  
 80 85 90

15  
 TCC CGC TCA TTG ACA CCC TGC ACC TGC GGC TCC TCG GAC CTT TAC 315  
 Ser Arg Ser Leu Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr  
 95 100 105

20  
 CTG GTT ACG AGG CAC GCC GAC GTC ATT CCC GTG CGC CGG CGA GGT 360  
 Leu Val Thr Arg His Ala Asp Val Ile Pro Val Arg Arg Arg Gly  
 110 115 120

25  
 GAT AGC AGG GGT AGC CTG CTT TCG CCC CGG CCC ATT TCC TAC CTA 405  
 Asp Ser Arg Gly Ser Leu Leu Ser Pro Arg Pro Ile Ser Tyr Leu  
 125 130 135

AAA GGC TCC TCG GGG GGT CCG CTG TTG TGC CCC GCG GGA CAC GCC 450  
 Lys Gly Ser Ser Gly Gly Pro Leu Leu Cys Pro Ala Gly His Ala  
 140 145 150

30  
 GTG GGC CTA TTC AGG GCC GCG GTG TGC ACC CGT GGA GTG ACC AAG 495  
 Val Gly Leu Phe Arg Ala Ala Val Cys Thr Arg Gly Val Thr Lys  
 155 160 165

35  
 GCG GTG GAC TTT ATC CCT GTG GAG AAC CTA GAG ACA ACC ATG AGA 540  
 Ala Val Asp Phe Ile Pro Val Glu Asn Leu Glu Thr Thr Met Arg  
 170 175 180

TCC CCG GTG

Ser Pro Val

## (2) INFORMATION FOR SEQ ID NO:2:

5

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 630 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

10

(D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: cDNA

## (ix) FEATURE:

15

(A) NAME/KEY: pT5His/HIV/183

ATG AGA GGA TCG CAT CAC CAT CAC CAT CAC GGA TCC CAT AAG GCA 45  
Met Arg Gly Ser His His His His His His Gly Ser His Lys Ala  
1 5 10 15

20

AGA GTT TTG GCT GAA GCA ATG AGC CAT GGT ACC ATG GCG CCC ATC 90  
Arg Val Leu Ala Glu Ala Met Ser His Gly Thr Met Ala Pro Ile  
20 25 30

25

ACG GCG TAC GCC CAG CAG ACG AGA GGC CTC CTA GGG TGT ATA ATC 135  
Thr Ala Tyr Ala Gln Gln Thr Arg Gly Leu Leu Gly Cys Ile Ile  
35 40 45

30

ACC AGC CTG ACT GGC CGG GAC AAA AAC CAA GTG GAG GGT GAG GTC 180  
Thr Ser Leu Thr Gly Arg Asp Lys Asn Gln Val Glu Gly Glu Val  
50 55 60

35

CAG ATC GTG TCA ACT GCT ACC CAA ACC TTC CTG GCA ACG TGC ATC 225  
Gln Ile Val Ser Thr Ala Thr Gln Thr Phe Leu Ala Thr Cys Ile  
65 70 75

AAT GGG GTA TGC TGG ACT GTC TAC CAC GGG GCC GGA ACG AGG ACC 270  
Asn Gly Val Cys Trp Thr Val Tyr His Gly Ala Gly Thr Arg Thr



- 25 -

	80	85	90
	ATC GCA TCA CCC AAG GGT CCT GTC ATC CAG ATG TAT ACC AAT GTG		315
5	Ile Ala Ser Pro Lys Gly Pro Val Ile Gln Met Tyr Thr Asn Val		
	95	100	105
	GAC CAA GAC CTT GTG GGC TGG CCC GCT CCT CAA GGT TCC CGC TCA		360
	Asp Gln Asp Leu Val Gly Trp Pro Ala Pro Gln Gly Ser Arg Ser		
10	110	115	120
	TTG ACA CCC TGC ACC TGC GGC TCC TCG GAC CTT TAC CTG GTT ACG		405
	Leu Thr Pro Cys Thr Cys Gly Ser Ser Asp Leu Tyr Leu Val Thr		
	125	130	135
15			
	AGG CAC GCC GAC GTC ATT CCC GTG CGC CGG CGA GGT GAT AGC AGG		450
	Arg His Ala Asp Val Ile Pro Val Arg Arg Arg Gly Asp Ser Arg		
	140	145	150
20			
	GGT AGC CTG CTT TCG CCC CGG CCC ATT TCC TAC CTA AAA GGC TCC		495
	Gly Ser Leu Leu Ser Pro Arg Pro Ile Ser Tyr Leu Lys Gly Ser		
	155	160	165
	TCG GGG GGT CCG CTG TTG TGC CCC GCG GGA CAC GCC GTG GGC CTA		540
25	Ser Gly Gly Pro Leu Leu Cys Pro Ala Gly His Ala Val Gly Leu		
	170	175	180
	TTC AGG GCC GCG GTG TGC ACC CGT GGA GTG ACC AAG GCG GTG GAC		585
	Phe Arg Ala Ala Val Cys Thr Arg Gly Val Thr Lys Ala Val Asp		
30	185	190	195
	TTT ATC CCT GTG GAG AAC CTA GAG ACA ACC ATG AGA TCC CCG GTG		630
	Phe Ile Pro Val Glu Asn Leu Glu Thr Thr Met Arg Ser Pro Val		
	200	205	210
35			

(2) INFORMATION FOR SEQ ID NO:3:

- 26 -

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: double

5 (D) TOPOLOGY: double

## (ii) MOLECULE TYPE: cDNA

10 GA TCA CCG GTC TAG ATCT  
T GGC CAG ATC TAGA

## 2) INFORMATION FOR SEQ ID NO:4:

15

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: polypeptide

## (ix) FEATURE:

25 (A) NAME/KEY: histidine tag

Met Arg Gly Ser His His His His His Thr Asp Pro

5

10

30

## (2) INFORMATION FOR SEQ ID NO:5:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 amino acids

35 (B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: polypeptide

(ix) FEATURE:

(A) NAME/KEY: Mutant Soluble 5A/5B Substrate

5

Asp Thr Glu Asp Val Val Ala Cys Ser Met Ser Tyr Thr Trp Thr  
5 10 15

Gly Lys

10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 162 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: cDNA

(ix) FEATURE:

(A) NAME/KEY: Native NS4A

25

TCA ACA TGG GTG CTC GTT GGC GGC GTC CTG GCT GCT CTG GCC GCG 45  
Ser Thr Trp Val Leu Val Gly Gly Val Leu Ala Ala Leu Ala Ala  
1 5 10 15

30

TAT TGC CTG TCA ACA GGC TGC GTG GTC ATA GTG GGC AGG ATT GTC 90  
Tyr Cys Leu Ser Thr Gly Cys Val Val Ile Val Gly Arg Ile Val  
20 25 30

TTG TCC GGG AAG CCG GCA ATT ATA CCT GAC AGG GAG GTT CTC TAC 135  
Leu Ser Gly Lys Pro Ala Ile Ile Pro Asp Arg Glu Val Leu Tyr  
35 40 45

35

CAG GAG TTC GAT GAG ATG GAA GAG TGC  
Gln Glu Phe Asp Glu Met Glu Glu Cys  
50

## (2) INFORMATION FOR SEQ ID NO:7:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 33 amino acid residues  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## 10 (ii) MOLECULE TYPE: polypeptide

## (ix) FEATURE:

- (A) NAME/KEY: Carboxl 33 mer of NS4A

15 Cys Val Val Ile Val Gly Arg Ile Val Leu Ser Gly Lys Pro Ala  
5 10 15  
Ile Ile Pro Asp Arg Glu Val Leu Tyr Gln Glu Phe Asp Glu Met  
20 25 30  
20 Glu Glu Cys

## (2) INFORMATION FOR SEQ ID NO:8:

25

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 amino acid residues  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
30 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: polypeptide

## (ix) FEATURE:

- 35 (A) NAME/KEY: Carboxl 33 mer of NS4A of HCV-BK strain

Ser Val Val Ile Val Gly Arg Ile Ile Leu Ser Gly Arg Pro Ala  
5 10 15

Ile Val Pro Asp Arg Glu Leu Leu Tyr Gln Glu Phe Asp Glu Met  
20 25 30

5 Glu Glu Cys

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- 10 (A) LENGTH: 20 amino acid residues  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: polypeptide

(ix) FEATURE:

(A) NAME/KEY: Soluble 4B/5A Substate

20 Trp Ile Ser Ser Glu Cys Thr Thr Pro Cys Ser Gly Ser Trp Leu  
5 10 15

Arg Asp Ile Trp Asp  
20

25

30

35

5 WE CLAIM:

1. A process for producing soluble, proteolytically active, refolded HCV protease from insoluble, bacterially produced HCV protease aggregates comprising:

10

(a) extracting insoluble, aggregates of HCV NS3 protease from bacteria producing said aggregates;

15

(b) solubilizing the aggregates of protease in a buffer containing a denaturing reagent;

(c) placing solubilized protease from step (b) in a buffer containing a reducing agent said buffer having an acidic pH;

20

(d) removing the denaturing reagent from the buffer under conditions wherein the buffer maintains an acidic pH; and

25

(e) raising the pH of the buffer containing protease in a stepwise manner to a pH of about 7 - 8 so as to produce properly refolded soluble, active NS3 protease.

2. The process of claim 1 wherein the denaturing agent is guanidine hydrochloride (GuHCl).

30

3. The process of claim 2 wherein the solution of GuHCl contains GuHCl at a concentration of about 5M.

4. The process of claim 1 wherein the reducing agent is dithiothreitol or  $\beta$ -mercaptoethanol.

35

5. The process of claim 1 wherein the buffer containing the reducing agent also contains a non-ionic detergent.



6. The process of claim 5 wherein the non-ionic detergent is selected from the group consisting of lauryl maltoside, a polyoxyethylene ether, nonidet P-40, a polyoxyethylene 9-lauryl ether such as, (3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate) (CHAPS),  
5 and octylglucoside.
7. The process of claim 6 wherein the solubilized protease of step (c) is in a 5M GuHCl solution and wherein the protease is reduced by diluting the 5M GuHCl solution with a buffer containing about 10mM DTT and  
10 0.1% lauryl maltoside.
8. The process of claim 1 wherein the denaturing reagent is removed in step (c) by applying the fractions containing the protease to a reverse phase chromatography column under conditions wherein fractions  
15 collected have an acidic pH.
9. The process of claim 8 wherein after the buffer containing the protease of step (c) is applied to the reverse phase chromatography column, a solution containing 99.9% H<sub>2</sub>O and 0.1% trifluoroacetic acid  
20 (TFA) is added to the column.
10. The process of claim 8 further comprising after adding the solution of 99.9% H<sub>2</sub>O + 0.1% TFA adding a solution comprised of 10% H<sub>2</sub>O + 90% acetonitrile + 0.1% TFA to the column at a 0 - 60% gradient into the  
25 solution of 99.9% H<sub>2</sub>O + 0.1% TFA and collecting the fractions.
11. The process of claim 10 further comprising dialyzing the fractions containing the properly refolded protein of step (c) first in an aqueous solution of 0.1% TFA resulting in a solution having a pH less than 1,  
30 then dialyzing the fraction in an aqueous solution of 0.01% TFA resulting in a solution having a pH of about 2 and then dialyzing the fractions in 0.001% TFA resulting in a solution having a pH of about 3, then dialyzing the solution in an aqueous solution having a pH of about 6, then dialysing the solution in an aqueous solution having a pH of  
35 about 7, then dialysing the solution in an aqueous solution having a pH of about 7.8 resulting in a solution having a pH of 7.4 - 7.8 containing properly refolded active HCV NS3 protease.

12. The process of claim 8 wherein after the buffer containing the protease of step (c) is applied to the reverse phase chromatography column, the column is eluted with a 0% - 60% acetonitrile gradient.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/06388

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 C07K1/113 //C12N15/51

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	FEBS LETT., vol. 324, 1993, pages 253-257, XP002013517 S. OSBORNE ET AL.: "Expression in E. coli and purification of a chimeric p22-NS3 recombinant antigen of Hepatitis C Virus (HCV)" see the whole document. ---	1
A	J. VIROL., vol. 67, 1993, pages 1385-1395, XP002013518 A. GRAKOU ET AL.: "Expression and indentification of hepatitis C virus polyprotein cleavage products" cited in the application see abstract and pages 1386-7. ---	1
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

24 September 1996

Date of mailing of the international search report

04. 10. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Yeats, S

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/06388

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 219 874 (BOEHRINGER MANNHEIM GMBH) 29 April 1987 see Examples 1a-1c and 2-5 and the claims. ---	1
A	EP,A,0 217 645 (CETUS CORPORATION) 8 April 1987 see Figure 1 and the claims. -----	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Int. Patent Application No

PCT/US 96/06388

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-219874	29-04-87	DE-A- 3537708	23-04-87
		AT-T- 131489	15-12-95
		AU-B- 607083	21-02-91
		AU-A- 4132189	04-01-90
		AU-B- 590029	26-10-89
		AU-A- 6599386	19-05-87
		CA-A- 1329157	03-05-94
		CZ-A- 8607526	17-01-96
		DE-D- 3650449	25-01-96
		DE-D- 3689404	27-01-94
		WO-A- 8702673	07-05-87
		EP-A- 0253823	27-01-88
		EP-A- 0393725	24-10-90
		ES-T- 2061434	16-12-94
		ES-T- 2020498	01-04-96
		FI-B- 94050	31-03-95
		FI-B- 95578	15-11-95
		HR-A- 921075	30-06-95
		IE-B- 62634	22-02-95
		JP-A- 4218387	07-08-92
		JP-B- 8024594	13-03-96
		JP-B- 7028745	05-04-95
		JP-T- 62502895	19-11-87
		US-A- 5453363	26-09-95
EP-A-217645	08-04-87	US-A- 4816440	28-03-89
		AU-B- 590618	09-11-89
		AU-A- 6318286	02-04-87
		CA-A- 1295245	04-02-92
		DE-A- 3681243	10-10-91
		JP-A- 62072625	03-04-87